

## Antioxidant and Anti-glycation Properties of a Combination of Seaweed and Mushroom and its Isolated Flavonoid over the Individual Extracts

Dhanush A, Badri Sai Madhulika, Usha C, Vaishnavi Musale, Pruthviraj K.S, Sinjitha .S. Nambiar\*

Department Of Chemistry of Biochemistry, School of Sciences, Jain (Deemed To Be) University, Bengaluru, Karnataka, India.

\*Corresponding author: [sinjithasn@gmail.com](mailto:sinjithasn@gmail.com)

Received: 24-05-2025; Accepted: 16-06-2025; Published: 28-06-2025

© Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License

<https://doi.org/10.55218/JASR.2025160606>

### ABSTRACT

Onset and progression of type 2 diabetes occurs due to oxidative stress and the formation of advanced glycation end products (AGEs). Dietary sources which contain natural antioxidants are being explored for their ability to combat these processes. In the current study, *Sargassum wightii*, a brown seaweed which is edible in nature and *Pleurotus djamor*, the pink oyster mushroom, were investigated individually and in combination for their antioxidant and anti-glycation activities. Water and ethanol extracts of the seaweed (SW) and mushroom (MS), as well as their mixtures (SW+MS), were prepared and evaluated using phosphomolybdate assay, reducing power assay, hydroxyl radical scavenging assay, and BSA-glucose antiglycation assay. The SW+MS water extract showed the highest total antioxidant activity (4.389 mg AAE/mg extract/ml) and a very high reducing power and hydroxyl radical scavenging capacity in comparison to individual extracts. Further, the flavonoid fraction isolated from the SW+MS combination showed the highest hydroxyl radical scavenging activity ( $IC_{50} = 0.8 \mu\text{g/ml}$ ), which was much higher than that of standard antioxidant. Moreover, the SW+MS ethanol extract showed the strongest anti-glycation potential among all tested samples. Our results show a synergistic interaction between *S. wightii* and *P. djamor*, thus providing a source of natural, antioxidant-rich food supplement with a therapeutic potential against diabetes.

**Keywords:** Antioxidant activity, Anti-glycation, *Pleurotus djamor*, *Sargassum wightii*, Seaweed-mushroom combination, Free radical scavenging, Flavonoids.

### INTRODUCTION

Free radicals cause formation of advanced glycation end products resulting in formation of sugar protein adducts which in turn lead to type 2 diabetes. These advanced glycation end products can in turn enhance free radical formation which in turn start a vicious cycle of advanced glycation end product formation [1]. To combat free radical formation, oxidative stress and its associated health risks, natural sources of antioxidants are being increasingly explored.

The edible brown algae *Sargassum wightii* has been known to have high antioxidant and antidiabetic activity [2,3]. *S. wightii* methanolic extracts have been shown to have antidiabetic potential in mice models of diabetes [3]. Seaweeds which are edible in nature show the presence of a large number of bioactive components like antioxidants, proteins, soluble dietary fibers, minerals, vitamins and polyunsaturated fatty acids [3]. Seaweeds have been used in traditional medicines in Asia against many diseases [3]. Studies have shown that seaweeds when consumed daily can prevent the occurrence of diseases like cardiovascular disease, cancers, hyperlipidemia etc. *Sargassum wightii*, seaweeds are present in Tamil Nadu, a state in India, abundantly. These are macroscopic, linear to ovate, with a height of 20–30 cm and length of 5–8 cm respectively [3].

Mushrooms have been known to provide protection against cancer and show antioxidant, anti-inflammatory, antitumor, antimicrobial activities. Mushrooms have been used as a source of natural food across the entire world being rich in protein, carbohydrates, crude fiber, vitamins, and minerals, as well as crude fat [4]. However, mushrooms have not been explored fully with respect to their therapeutic values. Mushrooms have also shown antibiotics and antioxidant properties [5].

*Pleurotus djamor*, pink oyster mushroom possesses excellent flavor, and is rich in nutrients [6,7,8]. Pink Oyster Mushroom (*Pleurotus Djamor*) is known to have antioxidant activity [9]. However, not much is known about its antidiabetic potential.

An imbalance between antioxidants and oxidants leads to the accumulation of free radicals (oxidative stress), which damage macromolecules like proteins, lipids, and nucleic acids. This damage can result in aging, abnormal gene expression, disruption of receptor activity, cell proliferation, immune perturbation, mutagenesis, tissue damage, and various disease conditions [10] like Alzheimer's disease, Parkinson's disease, muscular dystrophy, cataract, Rheumatoid Arthritis, diabetes, progeria, atherosclerosis, respiratory distress syndrome, Werner's syndrome, and ageing [11,10,12].

Seaweeds are known to contain excessive heavy metals which can adversely affect the body, if consumed in huge amounts [13]. Mushrooms have been shown to detoxify heavy metals from the body [14].

Hence, in this study, a combination of mushroom *Pleurotus djamor* and seaweeds *Sargassum wightii* has been explored for their antioxidant and anti-glycation potential to shed light on their future use as combinatorial food product which has anti-oxidant and anti-diabetic potential, but without the danger of experiencing heavy metal toxicity.

## MATERIALS AND METHODS

### Materials

- The pink oyster mushroom (*Pleurotus djamor*) was procured from, Green Aperon Ltd. Bangalore.
- The Brown seaweed algae (*Sargassum wightii*) was procured from Mandapam, Tamil Nadu.
- All the required chemicals and reagents were SRL labs, Bangalore.

### Methods

Water and ethanol extracts were prepared from *Sargassum wightii* (brown seaweeds) to make SW water and SW ethanol extracts. Water and ethanol extracts were prepared from *Pleurotus djamor* (pink oyster mushroom) to make MS water and MS ethanol extracts. Further a 1:1 combination of these extracts were prepared to make SW+MS water and SW+MS ethanol extracts

### Preparation of extracts

*Sargassum wightii* seaweed, *Pleurotus djamor* oyster mushroom alone or in ratio of 1:1 were dried overnight in a hot air oven overnight. The dried samples (*Sargassum wightii* seaweed, *Pleurotus djamor* oyster alone or in ratio of 1:1) were powdered. For preparing ethanol extracts, 50g of powdered samples were mixed with 50ml of 100% ethanol and homogenized and filtered using filter paper. The filtrates (of each sample) were collected in petriplates and left overnight for solvent evaporation. This results in preparation of *Sargassum wightii* seaweed (SW) ethanol extract, *Pleurotus djamor* oyster mushroom (MS) ethanol extract and *Sargassum wightii* seaweed, *Pleurotus djamor* oyster mushroom combination (SW+MS) ethanol extract.

For preparation of water extracts, 50g of powdered samples were mixed with 50ml of distilled water and homogenized and filtered using filter paper. The filtrates (of each sample) were lyophilized to obtain *Sargassum wightii* seaweed (SW) water extract, *Pleurotus djamor* oyster mushroom (MS) water extract and *Sargassum wightii* seaweed, *Pleurotus djamor* oyster mushroom combination (SW+MS) water extract.

### Isolation of flavonoids

Flavonoids were isolated from *Sargassum wightii* seaweed, *Pleurotus djamor* oyster mushroom using maceration method where the plants were ground in mortar and pestle containing 10g of sample to which 10-15ml of methanol is added. This is followed by filtration and drying of extracts [15].

### Total antioxidant assays (Phosphomolybdate assay)

Different concentrations of Ascorbic acid and 1mg/ml extracts were prepared. To each tube, ascorbic acid /extracts were taken and phosphomolybdate reagent ((70ml of reagent contained 0.23g of sodium phosphate monobasic and 0.33g of ammonium molybdate and 1.5 ml of concentrated Sulphuric acid) was added making the total reaction volume in each tube to be 2 ml. This was incubated in hot water bath (95°C) for 90 minutes. After 90min, the tubes are cooled to room temperature and their absorbances are read at 695nm. Results are expressed as mg ascorbic acid equivalent/ mg extract/ml of solution [16].

### Reducing power assay

Different aliquots of ascorbic acid and extracts of 0.002g/ml to 0.080g / ml were taken, to which phosphate buffer 0.2mM, pH 6.6, 0.1% of Ferric chloride and 1% of potassium ferricyanide were added and centrifuged at 6000rpm for 19 minutes. To the supernatant obtained after centrifugation, 10% TCA (trichloroacetic acid) was added and the solutions were incubated at room temperature for 30 minutes until – green colour appears. The absorbance is read at 700nm [17].

### Hydroxyl radical scavenging assay

The total reaction mixture (3.0ml) contained different aliquots of standard gallic acid/ extracts (2-10mg/ml), to which 1ml of ferrous -EDTA (0.13% of ferrous ammonium sulphate and 0.26% of EDTA) 0.5ml of EDTA, 1.0ml of chilled DMSO in phosphate buffer were added. To this reaction mixture 0.22% of ascorbic acid solution was added and incubated for 20 minutes. After cooling the solutions (in different tubes), ice cold 17.5% of TCA was added to all tubes. To this 1ml Nash reagent (0.075g of ammonium acetate, 0.03ml of glacial acetic acid, 0.02 ml of acetyl acetone in 50 ml water) was added and incubated at room temperature until the development of a pale yellow colour which was read at 412nm. Results were expressed as IC<sub>50</sub> (concentration of extract/standard causing 50% inhibition of hydroxyl radical production [18].

### Antiglycation assay

Different aliquots of extracts/standard were prepared and then 250 microlitre of 0.5mol/L glucose and 0.5mol/L Bovine serum albumin solutions are added into the reaction mixture, along with 0.2M pH 7.2 phosphate buffer with sodium azide. The solutions were incubated at 37°C for 3 days. After this, each reaction mixture was dialyzed against phosphate buffer, pH 7.2 for 24 h. After 24h, each of the solutions were mixed with 5% TCA & TBA and centrifuged at 3500rpm for 20 minutes. After this, the supernatants were collected in different tubes and absorbances were read at 412nm. Results were expressed as IC<sub>50</sub> (concentration of extract/standard causing 50% inhibition of glycation) [19, 20].

### Total phenolic content estimation.

Different concentrations are taken from stock gallic acid whose concentration was 1mg/ml. 50 microlitre of different concentrations of standard and 1mg/ml extract were added in different tubes and to these tubes, 65 microlitre of 8% sodium carbonate were added and

incubated for 6 minutes. Later FC reagent (1:1 ratio) was added to all tubes and incubated in dark for 1 hour. The blue colour developed was read at 765nm. Results were expressed in  $\mu\text{g}$  Gallic acid equivalent/ mg extract/ml of solution [21].

### Estimation of Total flavonoid content


Total flavonoid content was measured by preparing the following solutions: 0.5g of sodium nitrite in 10ml of distilled water, 10% Aluminium chloride solution and 0.1N sodium hydroxide. 10microlitres of different concentrations of standard flavonoid myricetin (stock 1mg/ml) and 1mg/ml extracts were taken in different tubes. 5 microlitres of sodium nitrite and then Aluminium chloride solution were added to the tubes and incubated for 10 minutes. To this 50 microlitres of sodium hydroxide solution was added and incubated for 5 minutes and absorbance was read at 510nm. Results were expressed in  $\mu\text{g}$  Myricetin equivalent/mg extract/ml of solution [21].

### Statistical Analysis

All the above experiments were performed in duplicates and statistically analyzed using ANOVA and Tukey test.


### RESULTS AND DISCUSSION

Seaweeds are rich in iodine, phenolic compounds, flavonoids, tannins, carotenoids and polysaccharides which enhance immunity and confer protection against free radicals [22]. *Pleurotus djamor* mushrooms were also shown to be rich in phenolic and flavonoid compounds which possess high antioxidant activities. However, the particular phytochemical responsible for its antioxidant activity has not been pinpointed and also the antioxidant activities of the combination of seaweeds and mushrooms remain unexplored till date. The present study focused on the phytochemical composition, antioxidant, anti-glycation, and free radical scavenging activities of different solvent extracts of seaweed (SW), mushroom (MS), and their mixture (SW+MS). The findings show that combining seaweed



SW- Seaweed, MS-Mushroom, SW+MS- mixture of seaweed and mushroom

**Fig. 1:** Total antioxidant activity (as assessed by phosphomolybdate assay) of seaweed-mushroom mixture water and seaweed-mushroom mixture ethanol extracts are compared with those of seaweed ethanol extract alone, seaweed water extract alone, mushroom ethanol extract alone, mushroom water extract alone and flavonoids isolated from seaweed-mushroom mixture. The results are expressed as mg ascorbic acid equivalent/ mg extract/ml. Each value represents mean  $\pm$  S.D. (n = 2).



SW- Seaweed, MS-Mushroom, SW+MS- mixture of seaweed and mushroom

**Fig. 2:** Reducing power assay. The reductive abilities of seaweed-mushroom mixture water and seaweed-mushroom mixture ethanol extracts at different doses. The reductive abilities are compared with those of seaweed ethanol extract alone, seaweed water extract alone, mushroom ethanol extract alone, mushroom water extract alone, flavonoids isolated from seaweed-mushroom mixture and standard Ascorbic acid (at different concentrations). The absorbance ( $A_{700}$ ) was plotted against the concentration of sample. Each value represents mean  $\pm$  S.D. (n = 2).

and mushroom increases bioactive compound content and associated biological activities.

### Antioxidant activity

Antioxidant activity was assessed using total antioxidant capacity (phosphomolybdate assay) and reducing power assays. The SW+MS water extract showed the highest total antioxidant activity (4.389 mg ascorbic acid equivalent/mg extract/ml), which was significantly higher than that of the SW+MS ethanol extract and extracts of SW or MS alone. Interestingly, the flavonoid fraction isolated from SW+MS showed equivalent antioxidant activity to the SW+MS water extract (Fig. 1), which further show that flavonoids are major contributors to antioxidant potential of SW+MS mixture.

SW- Seaweed, MS-Mushroom, SW+MS- mixture of seaweed and mushroom

The reducing power assay revealed that SW+MS water and ethanol extracts, as well as the flavonoid fraction extracted from the mixture, had higher reductive abilities than the individual extracts. The activity was found to be significantly higher than the standard antioxidant ascorbic acid (Fig. 2). The combinatorial interaction between SW and MS could potentially increase electron-donating capacity, which is important in neutralizing free radicals and preventing oxidative damage.

Previous studies have reported similar findings in seaweed and mushroom extracts, where phenolic compounds and flavonoids were identified as the primary contributors to antioxidant activity [23, 24]. The results from this study stress the role of flavonoids in participating in the antioxidant potential of the SW+MS mixture, making it a promising natural source of antioxidants.

### Free radical scavenging activity

Hydroxyl free radical starts off the free radical chain reaction which destroys lipids, proteins and nucleic acids [25]. Plants which scavenge hydroxyl free radical can be protective against damage to biomolecules.

Hydroxyl radical scavenging assay results showed the highly significant scavenging activities of SW+MS water and ethanol extracts (3.85 µg/ml and 3.9 µg/ml of IC<sub>50</sub> respectively) in comparison to individual SW and MS extracts whose activities were much lower than SW+MS extracts. The activities of SW+MS extracts were much higher than standard antioxidant mannitol with IC<sub>50</sub> = 200 µg/ml. Similar to the total antioxidant activity, the flavonoid fraction isolated from SW+MS showed the highest hydroxyl radical scavenging activity with IC<sub>50</sub> = 0.8 µg/ml, which was much higher than even the crude SW+MS extracts as well (Table 1).

This shows that the flavonoid fraction of the SW+MS mixture can scavenge hydroxyl radicals at extremely low concentrations and thus confer therapeutic activity to SW+MS mixture. Previously, it was seen that, the aqueous extract of *S. wightii* possesses hydroxyl radical scavenging activity. However, the bioactive compound responsible for its activity was not reported in the work [26]. Similarly, previously, mushroom extracts have been shown to have high hydroxyl radical scavenging activity [27]. However, the mixture of seaweeds and mushroom have never been explored for their combined antioxidant activities and compared with those of their individual extracts.

**Table 1:** Hydroxyl radical scavenging assay (in the form of IC<sub>50</sub>) for seaweed-mushroom mixture water and seaweed-mushroom mixture ethanol extracts, seaweed ethanol extract alone, seaweed water extract alone, mushroom ethanol extract alone, mushroom water extract alone, Flavonoids isolated from seaweed-mushroom mixture compared with the standard antioxidant Mannitol.

Extracts	IC <sub>50</sub> (µg/mL)
SW+MS water extract	3.85 ± 0.005 <sup>a</sup>
SW+MS ethanol extract	3.9 ± 0.000 <sup>a</sup>
MS water extract	80.5 ± 0.500 <sup>b</sup>
MS ethanol extract	72.5 ± 0.500 <sup>b</sup>
SW water extract	76.5 ± 0.500 <sup>b</sup>
SW ethanol extract	78.5 ± 0.500 <sup>b</sup>
Flavonoids from SW+MS	0.8 ± 0.000 <sup>c</sup>
Mannitol	200 ± 0.700 <sup>d</sup>

SW- Seaweed, MS-Mushroom, SW+MS- mixture of seaweed and mushroom  
The values shown here have been expressed as mean ± SD, *p* < 0.05. The experiments have been performed in duplicates. The letters given in superscript indicate the significance differences between the values after ANOVA and Tukey's test.

Note: Lower IC<sub>50</sub> value indicates higher activity.

### Anti-glycation activity

Glycation of globin proteins in hemoglobin is a major feature of diabetes. Plants which have anti-glycation activities confer protection against diabetes [28]. Advanced glycation end products (AGEs) are known to play a significant role in the pathogenesis of diabetes and associated complications. Among the ethanol extracts tested, SW+MS mixture had the highest anti-glycation activity IC<sub>50</sub> = 101 µg/ml compared with that of the positive anti-glycation standard rutin drug (IC<sub>50</sub> = 300 µg/ml). The SW+MS water extract also showed significant anti-glycation activity, with an IC<sub>50</sub> value of 125 µg/ml. The flavonoid fraction isolated from SW+MS mixture showed an IC<sub>50</sub> value of 140 µg/ml, indicating that flavonoids could be responsible for the significant antioxidant activity of the mixture. In comparison to individual SW and MS extracts, which possessed relatively weaker anti-glycation activity (IC<sub>50</sub> range of 208-278 µg/ml), SW+MS mixture showed a significant synergistic effect (Table 2). Thus, we can conclude that SW+MS extracts possess higher anti-glycation activity than individual extracts because their combined flavonoids can possibly scavenge more free radicals. The observed anti-glycation activity was consistent with previous studies where flavonoid compounds were found to inhibit protein glycation [29].

### Phytochemical analysis

The phytochemical screening of the extracts has shown a higher concentration of total flavonoid content (TFC) and total phenolic content (TPC) in SW+MS ethanol and water extracts than when each is extracted separately into SW or MS. Further, TFC and TPC of the water extracts of SW+MS mixture were found to be extremely high-about 2732.5 µg Myricetin equivalent/mg extract/ml and 2128.571 µg Gallic acid equivalent/mg extract/ml respectively (Table 3). These values were much greater (*p* < 0.05) than those of individual extracts, highlighting a synergistic effect that occurred upon combining SW and MS. The TFC isolated with the flavonoid-rich fraction of SW+MS

**Table 2:** Antiglycation activity (in the form of  $IC_{50}$ ) for seaweed-mushroom mixture water and seaweed-mushroom mixture ethanol extracts, seaweed ethanol extract alone, seaweed water extract alone, mushroom ethanol extract alone, mushroom water extract alone, Flavonoids isolated from seaweed-mushroom mixture compared with the standard anti-glycation drug Rutin.

Extracts	$IC_{50}$ ( $\mu\text{g}/\text{mL}$ )
SW+MS water extract	125 <sup>a</sup>
SW+MS ethanol extract	101 <sup>b</sup>
MS water extract	218 <sup>c</sup>
MS ethanol extract	210 <sup>c</sup>
SW water extract	208 <sup>c</sup>
SW ethanol extract	278 <sup>d</sup>
Flavonoids from SW+MS extract	140
Rutin	300 <sup>d</sup>

SW- Seaweed, MS-Mushroom, SW+MS- mixture of seaweed and mushroom  
The values shown here have been expressed as mean  $\pm$  SD,  $p < 0.05$ . The experiments have been performed in duplicates. The letters given in superscript indicate the significance differences between the values after ANOVA and Tukey's test.

Note: Lower  $IC_{50}$  value indicates higher activity

**Table 3:** Total flavonoid and total phenolic content of seaweed-mushroom mixture water and seaweed-mushroom mixture ethanol extracts, seaweed ethanol extract alone, seaweed water extract alone, mushroom ethanol extract alone, mushroom water extract alone and Flavonoids isolated from seaweed-mushroom mixture. Results have been expressed as microgram Myricetin equivalent/mg extract/ml for total flavonoid content (TFC) and as microgram Gallic acid equivalent/mg extract/ml for total phenolic content (TPC).

Extracts	TFC ( $\mu\text{g}$ myricetin equivalent / mg extract / ml)	TPC ( $\mu\text{g}$ gallic acid equivalent / mg extract / ml)
SW+MS water extract	2732.5 <sup>a</sup>	2128.571 <sup>b</sup>
SW+MS ethanol extract	2700 <sup>a</sup>	2142.857 <sup>b</sup>
MS water extract	2060 <sup>c</sup>	2007.143 <sup>d</sup>
MS ethanol extract	2095 <sup>c</sup>	2071.429 <sup>c</sup>
SW water extract	1060 <sup>e</sup>	1071.429 <sup>e</sup>
SW ethanol extract	1407.5 <sup>f</sup>	1378.571 <sup>f</sup>
Isolated flavonoid fraction from SW+MS	3571 <sup>g</sup>	

SW- Seaweed, MS-Mushroom, SW+MS- mixture of seaweed and mushroom  
The values shown here have been expressed as mean  $\pm$  SD,  $p < 0.05$ . The experiments have been performed in duplicates. The letters given in superscript indicate the significance differences between the values after ANOVA and Tukey's test.

was also the greatest observed (3571  $\mu\text{g}$  Myricetin equivalent/mg extract/ml), and serves as another confirmation that SW and MS are rich in flavonoids.

Flavonoids have been found to act as antioxidants by preventing free radical formation and also by scavenging free radical [30]. Their increased concentration in the SW+MS mixture could be potentially responsible for the superior biological activities observed in this study.

The values shown here have been expressed as mean  $\pm$  SD,  $p < 0.05$ . The experiments have been performed in duplicates. The letters given in superscript indicate the significance differences between the values after ANOVA and Tukey's test.

## CONCLUSION

The present study shows the synergistic effects of combining seaweed and mushroom, which highly enhance the phytochemical content and antioxidant activities. The SW+MS water and ethanol extracts showed superior antioxidant, anti-glycation, and hydroxyl radical scavenging activities compared to the individual extracts and the standard antioxidants. The isolated flavonoid fraction further showed excellent biological activity, thus indicating the importance of flavonoids as key bioactive compounds. The results indicate that mixtures of SW+MS, especially their aqueous and ethanol extracts, could prove to be powerful natural antioxidant and anti-glycation agents. The research presented here creates a very good foundation for further exploration of therapeutic effects of seaweed and mushroom combinations in the treatment of oxidative stress, diabetes, and related disorders. This should be followed by a focus on the isolation and structural characterization of active components and their in vivo evaluations to validate their efficacy.

## REFERENCES

- Ishrat N, Khan H, Patel O, Mahdi AA, Mujeeb F, Ahmad S. Role of Glycation in Type 2 Diabetes Mellitus and Its Prevention through *Nymphaea* Species. *BioMed Research International*. 2021; 7240046: 14 pages. Available from: <https://doi.org/10.1155/2021/7240046>.
- Vijayan R, Chitra L, Thiyagarajan R, Palvannan T. Dual antidiabetic and antihypertensive activity of fucoxanthin isolated from *Sargassum wightii* Greville in *in vivo* rat model. *Food Science and Human Wellness*. 2023; 12: 1693-1700. Available from: <https://doi.org/10.1016/j.fshw.2023.02.037>
- Emilin RR, Reenu N, Jane CPJ, Antony VS. Antidiabetic potential of methanolic extracts of *Sargassum wightii* in streptozotocin induced diabetic mice. *Biocatalysis and Agricultural Biotechnology*. 2020; 28: 101763. Available from: <https://doi.org/10.1016/j.bcab.2020.101763>.
- Boobalan M, Sethupathi N, Sengottuvelan PK, Balaji P, Gulyás B, Padmanabhan P, Selvan ST, Arun A. Mushroom-derived carbon dots for toxic metal ion detection and as antibacterial and anticancer agents. *ACS Applied Nano Materials*. 2020; 3: 5910-5919. Available from: <https://doi.org/10.1021/acsanm.0c01058>
- Ramanaiah I, Eyini M, Kumar M, Suresh Babu R, Prema P, Van-Huy Nguyen, Najat A. Bukhari, Ashraf A. Hatamleh, Balaji P. Bio-prospective potential of *Pleurotus djamor* and *Pleurotus florida* mycelial extracts towards Gram positive and Gram negative microbial pathogens causing infectious disease. *Journal of Infection and Public Health*. 2022; 15: 297-306. Available from: <https://doi.org/10.1016/j.jiph.2021.10.012>
- Dharmaraj K, Kuberan T, Mahalakshmi R. Comparison of nutrient contents and antimicrobial properties of *Pleurotus djamor*, *Agaricus bisporus* and *Ganoderma tsugae*. *International Journal of Current Microbiology and Applied Sciences*. 2014; 3: 518-26. Available from: ISSN: 2319-7706
- Lin SL, Lai CT, Ke X, Cheung PCK. Comparison of the composition and antioxidant activities of phenolics from the fruiting bodies of cultivated Asian culinary-medicinal mushrooms. *International Journal of Medicinal Mushrooms*. 2016; 18: 871-881. Available from: doi: 10.1615/intjmedmushrooms.v18.i10.30
- Hasan MT, Khatun MHA, Sajib MAM, Rahman MM, Rahman MS, Roy M. Effect of wheat bran supplement with sugarcane bagasse on growth, yield and proximate composition of pink oyster mushroom (*Pleurotus djamor*). *American Journal of Food Science*

- and Technology. 2015; 3:150–157. Available from doi: 10.12691/ajfst-3-6-2
9. Medeiros RL, Andrade GM, Crispim RB, Silva NNDS, Silva SAD, Souza HAN, Zárte-Salazar JR, Medeiros FD, Dantas CEA, Viera VB, Silva ALE, Tavares JF, Pereira FO. Nutritional and antioxidant potential of *Pleurotus djamor* (Rumph. ex Fr.) Boedijn produced on agronomic wastes banana leaves and sugarcane bagasse substrates. *Brazilian Journal of Microbiology*. 2024; 55:1117-1129. Available from: doi: 10.1007/s42770-024-01336-8.
  10. Martemucci G, Costagliola C, Mariano M, D'andrea L, Napolitano P, D'Alessandro AG. Free Radical Properties, Source and Targets, Antioxidant Consumption and Health. *Oxygen*. 2022; 2:48-78. Available from: <https://doi.org/10.3390/oxygen2020006>
  11. Vendemiale G, Grattagliano I, Altomare E. An update on the role of free radicals and antioxidant defense in human disease. *International Journal of Clinical and Laboratory Research*. 1999; 29: 49–55. Available from: <https://doi.org/10.1007/s005990050063>
  12. Jaeschke H, Gores GJ, Cederbaum AI, Hinson JA, Pessayre D, Lemasters JJ. Mechanisms of hepatotoxicity. *Toxicological sciences: an official journal of the Society of Toxicology*. 2002; 65: 166–176. Available from: <https://doi.org/10.1093/toxsci/65.2.166>.
  13. Lori H, Ji-Young L, Young-Ki P, Jaeun L. Heavy metals in seaweed: Implications for health benefits, risks, and safety regulations. *Journal of Agriculture and Food Research*. 2025; 21: 101830, Available from : <https://doi.org/10.1016/j.jafr.2025.101830>
  14. Zhang W, Zheng X, Chen X, Jiang X, Wang H, Zhang G. Lead detoxification of edible fungi *Auricularia auricula* and *Pleurotus ostreatus*: the purification of the chelation substances and their effects on rats. *Frontiers in Nutrition*. 2023; 10: 1162110. Available from: doi: 10.3389/fnut.2023.1162110.
  15. Tzanova M, Atanasov V, Yaneva Z, Ivanova D, Dinev T. Selectivity of Current Extraction Techniques for Flavonoids from Plant Materials. *Processes*. 2020; 8: 1222. Available from: <https://doi.org/10.3390/pr8101222>.
  16. Umamaheswari M, Chatterjee TK. In vitro antioxidant activities of the fractions of *Coccinia grandis* L. leaf extract. *African Journal of Traditional, Complementary and Alternative Medicines*. 2007; 5: 61–73. Available from: DOI:10.4314/ajtcam.v5i1.31258
  17. El Jemli M, Kamal R, Marmouzi I, Zerrouki A, Cherrah Y, Alaoui K. Radical-Scavenging Activity and Ferric Reducing Ability of *Juniperus thurifera* (L.), *J. oxycedrus* (L.), *J. phoenicea* (L.) and *Tetraclinis articulata* (L.). *Advances in Pharmacological Sciences*. 2016; 2016: 6392656. Available from doi: 10.1155/2016/6392656.
  18. Suseela V, Sushmita L, Bharatkumar R, Nirmaladevi R. Free Radical Scavenging potential of different extracts of *Tabebuia roseo-alba* (Ridl) Sand leaves. *Research Journal of Pharmacy and Technology*. 2021; 14: 4801-7. Available from doi: 10.52711/0974-360X.2021.00835.
  19. Kennedy DM, Skillen AW, Self CH. Colorimetric assay of glycoprotein glycation free of interference from glycosylation residues. *Clinical chemistry*, 1993; 39: 2309–2311. Available from PMID: 7693374
  20. Halliwell B, Gutteridge JM, Aruoma, OI. The deoxyribose method: a simple “test-tube” assay for determination of rate constants for reactions of hydroxyl radicals. *Analytical biochemistry*, 1987; 165: 215–219. Available from: [https://doi.org/10.1016/0003-2697\(87\)90222-3](https://doi.org/10.1016/0003-2697(87)90222-3).
  21. Samidha K, Vrushali K, Vijaya P. Estimation of Phenolic content, Flavonoid content, Antioxidant and Alpha amylase Inhibitory Activity of Marketed Polyherbal Formulation, *Journal of Applied Pharmaceutical Science*. 2014; 4: 061-065. Available from: DOI 10.7324/JAPS.2014.40911.
  22. Michalak I, Tiwari R, Dhawan M, Alagawany M, Farag MR, Sharun K, Emran TB, Dhama K. Antioxidant effects of seaweeds and their active compounds on animal health and production - a review. *The Veterinary Quarterly*. 2022; 42: 48-67. Available from doi: 10.1080/01652176.2022.2061744. PMID: 35363108; PMCID: PMC9004519.
  23. Cotas J, Leandro A, Monteiro P, Pacheco D, Figueirinha A, Gonçalves AMM, da Silva GJ, Pereira L. Seaweed Phenolics: From Extraction to Applications. *Marine Drugs*. 2020; 18: 384. Available from doi: 10.3390/md18080384.
  24. Azieana J, Zainon M, Noriham A, Rohana M. Total Phenolic and Flavonoid Content and Antioxidant Activities of Ten Malaysian Wild Mushrooms. *Open Access Library Journal*, 2017; 4: 1-9. Available from doi: 10.4236/oalib.1103987.
  25. Ronald LP. Oxygen radical absorbance capacity (ORAC): New horizons in relating dietary antioxidants/bioactives and health benefits, *Journal of Functional Foods*, 2015; 18: 797-810. Available from: <https://doi.org/10.1016/j.jff.2014.12.018>.
  26. Sradhasini R, Bandana R, Subrat KB, Ishani R, Anjan K, Antioxidant and anti-inflammatory activities of methanol and aqueous extracts of *Sargassum wightii*. *Journal of Herbmed Pharmacology*. 2022; 11: 75-82. Available from: doi: 10.34172/jhp.2022.08
  27. Muna G, John M, Benson M, Ogoyi D. African Journal of Biotechnology Antioxidant properties of cultivated edible mushroom (*Agaricus bisporus*) in Kenya. *African Journal of Biotechnology*. 2015; 14: 1401-1408. Available from: doi: 10.5897/AJB2015.14436.
  28. Mahfuza A, Faria FP, Noore ZN, Nasim A, Md. Aatur R, Md. Masum B, Md. Abdul B, Sabbir A, Md. Nurul I. Antioxidant, anti-glycation and hypoglycemic potentials of *Acmella uliginosa*: A probable candidate for the amelioration of diabetes mellitus. *Phytomedicine Plus*. 2025; 5: 100777, Available from: doi: <https://doi.org/10.1016/j.phyplu.2025.100777>.
  29. Patil KK, Meshram RJ, Barage SH, Gacche RN. Dietary flavonoids inhibit the glycation of lens proteins: implications in the management of diabetic cataract. *3 Biotech*. 2019; 9: 47. Available from doi: 10.1007/s13205-019-1581-3.
  30. Pietta PG. Flavonoids as antioxidants. *Journal of Natural Products*. 2000; 63: 1035-42. Available from doi: 10.1021/np9904509.

**HOW TO CITE THIS ARTICLE:** Dhanush A, Madhulika BS, Usha C, Musale V, Pruthviraj KS, Nambiar SS. Antioxidant and Anti-glycation Properties of a Combination of Seaweed and Mushroom and its Isolated Flavonoid over the Individual Extracts. *J Adv Sci Res*. 2025; 16(06): 56-61 DOI: 10.55218/JASR.2025160606